Determining the Mechanisms Limiting Amphibian Distributions in Urban Areas

> Katja Kwaku CAWS Annual Meeting March 9, 2023



Species have varying sensitivities to urbanization, likely depending on:

- Hydroperiod requirements
- Dispersal ability
- Habitat flexibility



Adapted from Gibbs 1998

Focal Cities



Milford (CT)

Hartford (CT)

Cambridge (MA)

Urban	Rural	Urban	Rural	Urban	Rural
Milford	Orange	Hartford, Eas	t West Hartford	Cambridge,	Weston, Wayland
		Hartford		Arlington	
• Spring peepers	• Spring peepers	• Green frogs	• Spring peepers	 Bullfrogs 	Spring peepers
	Wood frogs	American to	ads • Wood frogs		Wood frogs
			Spotted		 Spotted
			salamanders		salamanders
			• Gray tree frogs		Green frogs
			Green frogs		American toads



Predicted Results

Wood frogs

Rural

Spring peepers

Problem: Poor terrestrial habitat quality



Survival Rate

Problem: Dispersal limitation



Urban

Rural

Urban



Conservation Implications



Thanks for listening!

Acknowledgements

- Advisor: Dr. Mark C. Urban
- Field Assistants: Finn O'Shaughnessey, Carmella Testa, Geneva Kwaku, Ketki Samel



Michael Lefor Grant, CAWS



Biology Graduate Student Research Endowment Fund to the Department of Ecology and Evolutionary Biology



This material is based upon work supported in part by the National Science Foundation under Grant DGE-2022036. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the National Science Foundation.